

When Newton meets Descartes: A Simple and Fast Algorithm to Isolate the Real Roots of a Polynomial

Michael Sagraloff

MPI for Informatics, Saarbrücken, Germany

Abstract

We introduce a new algorithm denoted DSC^2 to isolate the real roots of a univariate square-free polynomial f with integer coefficients. The algorithm iteratively subdivides an initial interval which is known to contain all real roots of f . The main novelty of our approach is that we combine Descartes' Rule of Signs and Newton iteration. More precisely, instead of using a fixed subdivision strategy such as bisection in each iteration, a Newton step based on the number of sign variations for an actual interval is considered, and, only if the Newton step fails, we fall back to bisection. Following this approach, our analysis shows that, for most iterations, we can achieve quadratic convergence towards the real roots. In terms of complexity, our method induces a recursion tree of almost optimal size $O(n \cdot \log(n\tau))$, where n denotes the degree of the polynomial and τ the bitsize of its coefficients. The latter bound constitutes an improvement by a factor of τ upon all existing subdivision methods for the task of isolating the real roots. In addition, we provide a bit complexity analysis showing that DSC^2 needs only $\tilde{O}(n^3\tau)$ bit operations¹ to isolate all real roots of f . This matches the best bound known for this fundamental problem. However, in comparison to the much more involved algorithms by Pan and Schönhage (for the task of isolating all complex roots) which achieve the same bit complexity, DSC^2 focuses on real root isolation, is very easy to access and easy to implement.

Key words: Root isolation, Subdivision Methods, Exact Computation, Newton's Method, Descartes' Rule of Signs, Bit Complexity, Asymptotically Fast Methods

1. Introduction

Finding the roots of a univariate polynomial f is considered as one of the most important tasks in computational algebra. This is justified by the fact that many problems from mathematics, engineering, computer science, and the natural sciences can be reduced to solving a system of polynomial equations which in turn, by means of elimination techniques such as resultants or Gröbner Bases, reduces to solving a polynomial equation in one variable. Hence, it is not

¹ $\tilde{O}$ indicates that we omit logarithmic factors.

Email address: msagralo@mpi-inf.mpg.de (Michael Sagraloff).

surprising that numerous approaches are dedicated to this fundamental problem. We mainly distinguish between (1) numerical and exact methods, and (2) methods to find all complex roots and methods which are especially tuned to search for real roots. The numerical literature lists many algorithms, such as Newton iteration or the Weierstrass-Durand-Kerner method, that are widely used and effective in practice but lack a guarantee on the global behavior (cf. [24] for discussion). In particular, the convergence and/or the complexity of the Weierstrass-Durand-Kerner method is still open.

The work of A. Schönhage [29] from 1982 marks the beginning of the complexity-theoretic approaches. It combines a newly introduced concept denoted *splitting circle method* with techniques from numerical analysis (Newton iteration, Graeffe's method, discrete Fourier transforms) and fast algorithms for polynomial and integer multiplication. For the benchmark problem of isolating all complex roots of a polynomial f of degree n with integer coefficients of modulus 2^τ or less, the proposed method achieves the record bound of $\tilde{O}(n^3\tau)$ bit operations. V. Pan and others [23,24] gave theoretical improvements in the sense of achieving record bounds simultaneously in both bit complexity and arithmetic complexity, but the initial bound $\tilde{O}(n^3\tau)$ on the number of bit operations has still remained intact. Common to all asymptotically fast algorithms is (as the authors themselves admit) that they are rather involved and very difficult to implement. The latter is also due to the fact that, in order to control the precision errors in the considered numerical subroutines, one has to carefully work out many details of their implementation. Hence, it is not surprising that, despite their theoretical richness, the asymptotically fast algorithms have so far not been used, or not proven to be efficient in practice; see [12] for an implementation of the splitting circle method within the Computer Algebra system Pari/GP. A further reason might be that the benchmark problem is inappropriate for most applications. For instance, in ray shooting in computer graphics, we are only interested in the first positive root or in the real roots in some specified neighborhood.

In parallel to the development of purely numerical methods, there is a steady ongoing research on exact subdivision algorithms, such as the Descartes method [7,9,15,18,26], the Bolzano method [4,5,6,17,33], Sturm Sequences [8,16,25] or the continued fraction method [2,30,31]. These methods from the exact computation literature are widely used in various algebraic applications (e.g., cylindrical algebraic decomposition), and many of them have been integrated into computer algebra systems (e.g., MAPLE, MATHEMATICA, SAGE, etc.). In addition, their computational complexity has been well-studied [8,10,30,31,33], and many experiments have shown their practical evidence [13,14,26]. Current experimental data shows that a version of the Descartes method (i.e., the univariate solver in RS based on [26], integrated into MAPLE) which uses approximate computation performs best for most polynomials, whereas, for harder instances, the continued fraction approach seems to be more efficient. With respect to the benchmark problem, all of the above mentioned algorithms demand for $\tilde{O}(n^4\tau^2)$ bit operations to isolate all real roots, hence they tend to lag behind the asymptotically fast algorithms by a factor of $n\tau$. Recently, it has been shown [28] that the bound on the bit complexity for the Descartes method can be lowered to $\tilde{O}(n^3\tau^2)$ when replacing exact computation by approximate computation (without abstaining from correctness). This result partially explains the success of such a modified Descartes method in practice. However, as long as we restrict to the bisection strategy, it seems that the latter bound is optimal. We remark that Schönhage already made a similar observation: In the introduction of [29], he argued that "a factor τ^2 inevitably occurs if nothing better than linear convergence is achieved".

In this paper, we introduce an exact and complete subdivision algorithm denoted DSC^2 to isolate all real roots of a square-free polynomial with integer coefficients. Similar to the classical Descartes method, we use Descartes' Rule of Signs to determine an upper bound $v_I = \text{var}(f, I)$ for the number of real roots of the polynomial f within an interval I that is actually processed. However, instead of splitting I into two equally sized subintervals in each iteration, we consider a subdivision strategy that is based on Newton iteration: The analysis of the classical approach which exclusively uses bisection shows that the induced recursion tree is large ($\approx n\tau$) if and only if there exists a long sequences $I_1 \supset I_2 \supset \dots \supset I_s$ of intervals in the subdivision process, where $v = v_{I_1} = \dots = v_{I_s}$. Such a sequence implies the existence of a cluster $\mathcal{C}$ of v nearby roots, and vice versa (cf. Theorem 1). Hence, it seems reasonable to obtain a good approximation (i.e., an interval $I' \subset I$ close to $\mathcal{C}$) of such a cluster by considering a corresponding Newton step to approximate a v -fold root. Combining Descartes' Rule of Signs and a subdivision technique similar to the one as proposed by J. Abbott for the QIR method [1], we formulate a method to determine whether the so-obtained approximation I' should be kept or not. In case of success, we proceed with the considerably smaller interval I' , whereas we fall back to bisection if the Newton step fails. Our analysis shows that, following this approach, we achieve quadratic convergence in most iterations. As a consequence, the induced recursion tree has almost optimal size $O(n \log(n\tau))$ which improves upon the bisection strategy by a factor of τ . We further provide a detailed bit complexity analysis which yields the bound $\tilde{O}(n^3 \tau)$ for DSC^2 . This matches the record bound achieved by the aforementioned asymptotically fast algorithms.

We consider our contribution of great importance because of the following reasons: (1) Although the proposed method is rather simple, it achieves the best bounds known for the bit complexity of the problem of isolating the real roots of a polynomial. (2) In addition, it is much easier to access and also much easier to implement than the asymptotically fast algorithms that are available so far. In comparison to the existing practical methods for real root isolation, the modifications are moderate, and thus we expect that a careful implementation of our new approach will outperform the existing ones. (3) Finally, our method can be applied to search for the real roots in some specified neighborhood of interest, a property which is not fulfilled by the algorithms as proposed by Pan and Schönhage.

2. Overall Idea

In this section, we first provide a high-level description of our new algorithm, and then outline the argument why this approach improves upon existing methods such as the classical Descartes method. For the exact definition of the algorithm and a detailed bit complexity analysis, we refer the interested reader to Section 3. Throughout the following considerations, let

$$f(x) := \sum_{i=0}^n a_i x^i \in \mathbb{Z}[x], \text{ with } |a_i| < 2^\tau, \quad (2.1)$$

be a square-free polynomial of degree n with integer coefficients of bit-length τ or less. We further denote $z_1, \dots, z_n$ the complex roots of f , $\sigma(z_i) := \min_{j \neq i} |z_i - z_j|$ the *separation of z_i* , and $\sigma_f := \min_i \sigma(z_i)$ the *separation of f* . According to Cauchy's bound, the modulus of each root z_i is bounded by $1 + 2^\tau \leq 2^{\tau+1}$, and thus, for the task of isolating the real roots of f , we can restrict our search to the interval $\mathcal{I}_0 := (-2^{\tau+1}, 2^{\tau+1})$.

We consider an arbitrary root isolation method denoted ISO which recursively performs subdivision on I_0 in order to determine isolating intervals for the real roots of f . ISO uses a counting




Fig. 2.1. For any k with $0 \leq k \leq n$, the Obreshkoff discs $\bar{C}_k$ and $\underline{C}_k$ for $I = (a, b)$ have the endpoints of I on their boundaries; their centers see the line segment (a, b) under the angle $2\pi/(k+2)$. The Obreshkoff lens L_k is the interior of $\bar{C}_k \cap \underline{C}_k$, and the Obreshkoff area A_k is the interior of $\bar{C}_k \cup \underline{C}_k$. Any point (except for a and b) on the boundary of A_k sees I under an angle $\pi/(k+2)$, and any point (except for a and b) on the boundary of L_k sees I under the angle $\pi - \pi/(k+2)$. We have $L_n \subset \dots \subset L_1 \subset L_0$ and $A_0 \subset A_1 \subset \dots \subset A_n$. The cases $k = 0$ and $k = 1$ are of special interest: The circles $\bar{C}_0$ and $\underline{C}_0$ coincide. They have their centers at the midpoint of I . The circles $\bar{C}_1$ and $\underline{C}_1$ are the circumcircles of the two equilateral triangles having I as one of their edges. We call A_0 and A_1 the *one* and *two-circle regions* for I .

function $\text{var}(f, I)$ for m , the number of roots within an interval $I \subset \mathcal{I}_0$, where $v := \text{var}(f, I) \in \mathbb{N}$ fulfills the following properties:

- (P1) v is an upper bound for m (i.e., $v \geq m$), and
- (P2) v has the same parity as m (i.e., $v \equiv m \pmod{2}$).

The latter two properties imply that $\text{var}(f, I) = m$ if $v \leq 1$. Hence, in each step of the recursion, an interval I is stored as isolating if $v = 1$, and I is discarded if $v = 0$. If $v > 1$, $I = (a, b)$ is subdivided (according to some subdivision strategy) into subintervals $I_1 = (a, \lambda_1)$, $I_2 = (\lambda_1, \lambda_2)$, $\dots$, $I_l := (\lambda_{l-1}, b)$, where $1 \leq l \leq l_0$, $l_0 \in \mathbb{N}$ is a global constant, and the λ_i are rational values. In order to detect roots at the subdivision points λ_i , we also check whether $f(\lambda_i) = 0$ or not. T_{ISO} denotes the recursion tree induced by ISO. Throughout the following considerations, we often treat nodes of ISO and intervals produced by ISO as interchangeable.

The definition of the counting function $\text{var}(f, I)$ is based on Descartes' Rule of Signs: *For an arbitrary polynomial $p = \sum_{i=0}^n p_i x^i \in \mathbb{R}[x]$, the number m of positive real roots of p is bounded by the number v of sign variations in its coefficient sequence $(p_0, \dots, p_n)$ and, in addition, $v \equiv m \pmod{2}$.* In order to extend the latter rule to arbitrary intervals $I = (a, b)$, the Möbius transformation $x \mapsto \frac{ax+b}{x+1}$ which maps $(0, +\infty)$ one-to-one onto I is considered. Thus, for

$$f_I(x) = \sum_{i=0}^n c_i x^i := (x+1)^n \cdot f\left(\frac{ax+b}{x+1}\right), \quad (2.2)$$

and $\text{var}(f, I)$ defined as the number of sign variations in the coefficient sequence $(c_0, \dots, c_n)$ of f_I , $\text{var}(f, I)$ fulfills the properties (P1) and (P2). Because of the latter two properties and the fact that we never discard intervals that contain a real root of f , correctness of ISO follows immediately.

The *classical Descartes method* (DSC for short) is a subdivision method which uses bisection in each iteration, that is, in each step, we have $l = l_0 = 2$, $I_1 = (a, \lambda_1) := (a, m(I))$ and

$I_2 = (\lambda_1, b) = (m(I), b)$, with $m(I) := (a+b)/2$ the midpoint of I . Termination and complexity analysis of DSC rest on the following theorem:

Theorem 1 ([19,21]). *Let $I = (a, b)$ be an open interval and $v = \text{var}(f, I)$. If the Obreshkoff lens $L_{n-k} \subset \mathbb{C}$ (see Figure 2.1 for the definition of L_{n-k}) contains at least k roots (counted with multiplicity) of f , then $v \geq k$. If the Obreshkoff area $A_k \subset \mathbb{C}$ contains at most k roots (counted with multiplicity) of k , then $v \leq k$. In particular,*

$$(P3) \quad \# \text{ of roots of } f \text{ in } L_n \leq \text{var}(p, I) \leq \# \text{ of roots of } f \text{ in } A_n.$$

We remark that the special cases $k = 0$ and $k = 1$ appear as the one- and two-circle theorems in the literature [3,9,15,20,22]. For the Descartes method, Theorem 1 implies that no interval I of length $w(I) \leq \sigma_f$ is split. Namely, its one-circle region A_0 cannot contain two or more roots. If A_0 contains no root, then $\text{var}(f, I) = 0$. Otherwise, A_0 contains one real root, and thus the two-circle region A_1 contains no non-real root. Hence, in the latter situation, we have $\text{var}(f, I) = 1$ by Theorem 1. We conclude that the depth of the recursion tree T_{DSC} induced by the Descartes method is bounded by $\log w(\mathcal{J}_0) + \log \sigma_f^{-1} = \tau + \log \sigma_f^{-1} + 2$. Furthermore, it holds (see [9, Corollary 2.27] for a self-contained proof):

Theorem 2. *Let I be an interval and I_1 and I_2 be two disjoint subintervals of I . Then,*

$$(P4) \quad \text{var}(f, I_1) + \text{var}(f, I_2) \leq \text{var}(f, I).$$

According to Theorem 2, there cannot be more than $n/2$ intervals I with $\text{var}(f, I) \geq 2$ at any level of the recursion. Hence, the size of T_{DSC} is bounded by $n(\tau + \log \sigma_f^{-1} + 2)$. Using Davenport-Mahler bound, one can further show [9,28] that $\log \sigma_f^{-1} = O(n(\log n + \tau))$, and thus the bound for $|T_{\text{DSC}}|$ writes as $\tilde{O}(n^2\tau)$. A more refined argument [9] yields $|T_{\text{DSC}}| = \tilde{O}(n\tau)$ which is optimal for the bisection strategy.

In the next step, we study the situation where the recursion tree T_{DSC} for the classical Descartes method is large. We then introduce our new algorithm which we denote DSC^2 due to the quadratic convergence in most steps. DSC^2 is a variant of the Descartes method which adaptively addresses the latter situation via combining Newton iteration and bisection. We also sketch the argument why this approach improves upon DSC. The following definition is essential for the argument; see also Figure 2.2:

Definition 1. Let I be a node (interval) in the recursion tree T_{ISO} induced by some subdivision algorithm ISO. We call I *terminal* if $\text{var}(f, I) \leq 1$. A non-terminal interval I with children $I_1, \dots, I_l$ is called *special* if $I = \mathcal{J}_0$ (i.e., I is the root of T_{ISO}), or

$$\text{var}(f, I_j) < \text{var}(f, I) \text{ for all } j = 1, \dots, s.$$

According to (P4) in Theorem 2, we have $\sum_{j=1}^l \text{var}(f, I_j) \leq \text{var}(f, I)$ for each I . Thus, a non-terminal node I different from $\mathcal{J}_0$ is non-special if and only if, for one of its children, we count the same number of sign variations as for I and, for all other children, we count no sign variation. In total, there exist n' special nodes, where $n' \leq \text{var}(f, \mathcal{J}_0) \leq n$. Namely, when we subdivide a special interval which is not the root of the recursion tree, the non-negative value $\mu := \sum_I \text{var}(f, I) - \#\{I : \text{var}(f, I) > 0\}$ decreases by at least one, where we sum over all leafs in the actual iteration, and μ is initially set to $\mu = \text{var}(f, \mathcal{J}_0) - 1 \leq n - 1$. We denote the special




Fig. 2.3. The long chain $I_1 \supset I_2 \supset \dots \supset I_s$ of intervals in $T(5)$ with $\text{var}(f, I_1) = \dots = \text{var}(f, I_s) = 2$ corresponds to a large number of bisection steps to isolate two very nearby roots from each other; see the figure on the right with the graph of f over I_1 .

to (P3) in Theorem 1, there exists a long sequence $I_1 \supset I_2 \supset \dots \supset I_s$ of non-special intervals with $\xi \in I_j$ for all j , and thus the number

$$v := \text{var}(f, I_1) = \text{var}(f, I_2) = \dots = \text{var}(f, I_s)$$

of sign variations does not change for the intervals in this sequence. Namely, for each I_j in the above sequence, the Obreshkoff lens L_n contains $\mathcal{C}$. Vice versa, according to Theorem 1, such a long sequence of non-special intervals implies the existence of a cluster $\mathcal{C}$ consisting of v nearby roots as above because the Obreshkoff area A_n of each I_j must contain at least v roots.² Since a cluster $\mathcal{C}$ of v nearby roots at ξ behaves very similar to a v -fold root at ξ , it seems reasonable to obtain a good approximation of $\mathcal{C}$ by considering Newton iteration instead of bisection. Namely, for a polynomial $p(x) \in \mathbb{R}[x]$ with a v -fold root at ξ and a starting value x_0 sufficiently close to ξ , it is well-known from numerical analysis that the sequence $(x_i)_{i \in \mathbb{N}_0}$ recursively defined by

$$x_{i+1} := x_i - v \cdot \frac{p(x_i)}{p'(x_i)}$$

converges quadratically to ξ . Unfortunately, when isolating the roots of f , the situation differs considerably from the latter one: First, the above result only holds for a v -fold root ξ and does not directly extend to a cluster $\mathcal{C}$ of v roots near ξ . Second, in an early stage of the subdivision process, the existence of such a cluster $\mathcal{C}$ is not guaranteed, and even if one exists, we do not know what "sufficiently close to ξ " means in this situation.

² The thoughtful reader may notice that the latter two statements are not completely rigorous: In particular, in the special case where ξ , and thus also the cluster $\mathcal{C}$, is very close to one of the endpoints of some I_j , it might happen that some of the roots are not considered by the counting function $\text{var}(f, I_j)$ since they are located outside the Obreshkoff lens/area. We will address this issue in our algorithm as defined in Section 3.

In order to address the above mentioned problems and to finally turn the purely numerical Newton method into an exact and complete algorithm, we propose the following approach: Let $v = \text{var}(f, I)$ be the number of sign variation for an actual interval $I = (a, b)$ in a certain iteration. Then, we consider this as an indicator that there might exist a cluster of v nearby roots. Thus, we compute $\lambda := t - v \cdot f(t)/f'(t)$ for some $t \in [a, b]$ (e.g., an endpoint of I) and consider an interval $I' = (a', b') \subset I$ of width $w(I') \ll w(I)$ that contains λ . If $\text{var}(f, I') = v$ as well, we keep I' and discard the intervals $(a, a']$ and $[b', b)$. Otherwise, we split I into two equally sized intervals $I_1 := (a, m(I))$ and $I_2 := (m(I), b)$ and finally check whether $f(m(I)) = 0$ or not. Following this approach, no root is lost and intervals are at least bisected in each iteration. Furthermore, if a cluster $\mathcal{C}$ of nearby roots actually exists, we can hope to achieve fast convergence to this cluster when choosing I' in an appropriate manner. In our algorithm, we choose I' in a similar way as proposed by Abbott [1] for the task of further refining intervals which are already isolating for an ordinary root. Namely, we decompose I into a certain number N_I of subintervals and pick the subinterval I' of size $w(I)/N_I$ which contains λ . If $\text{var}(f, I') = v$, then we keep I' and decompose I' into $N_{I'} = N_I^2$ subintervals in the next iteration. Otherwise, we continue with the intervals $I_1 = (a, m(I))$ and $I_2 = (m(I), b)$ which are now decomposed into only $N_{I_1} = N_{I_2} := \max(4, \sqrt{N_I})$ many subintervals, etc.

In the next section, we give the exact definition of our new algorithm, and we show that it induces a subdivision tree of considerably smaller size than T_{DSC} . In particular, it turns out that the size of each $T_i \subset T_{\text{DSC}^2}$ is bounded by $O(\log n + \log \tau)$ which is due to the fact that, for most iterations, we have quadratic convergence to the real roots, and the width of each interval is lower bounded by $2^{-\tilde{O}(n\tau)}$; see Lemma 1 and Theorem 4 for proofs. Hence, according to (2.3), the size of the overall recursion tree is bounded by

$$O(n' \cdot (\log n\tau)) = O(\text{var}(f, \mathcal{J}_0) \cdot (\log n\tau)) = O(n \cdot (\log n\tau)). \quad (2.4)$$

The latter result particularly shows that the size of the recursion tree is directly correlated to the number n^* of non-zero coefficients of f because instead of considering $\mathcal{J}_0 = (-2^{\tau+1}, 2^{\tau+1})$, we can start with $(-2^{\tau+1}, 0)$ and $(0, 2^{\tau+1})$, and the total number of sign variations counted for both intervals is upper bounded by $2 \cdot n^*$.

For the bit complexity of our algorithm, we have to consider the costs for computing the polynomials $f_I(x) = (1+x)^n \cdot f((ax+b)/(1+x))$ as defined in (2.2), where $I = (a, b)$ is an interval to be processed. In Section 3.3, we will show that the costs for the latter step are dominated (up to constant factors) by the computation of $f(x+a)$. For $I \in T_i$, the endpoints of I are dyadic numbers of bitsize $O(\tau + \log w(J_i)^{-1})$ or less, and thus the computation of f_I demands for $\tilde{O}(n^2(\tau + \log w(J_i)^{-1}))$ bit operations. In Lemma 6, we prove that we can order the roots $z_1, \dots, z_n$ in a way such that $\log w(J_i)^{-1} < \log \sigma(z_i)^{-1} + O(\log^2 n)$ for all $i = 1, \dots, n'$. Then, it follows that computing f_I demands for at most

$$\tilde{O}(n^2(\tau + \log w(J_i)^{-1})) = \tilde{O}(n^2(\tau + \log \sigma(z_i)^{-1} + \log^2 n)) = \tilde{O}(n^2(\tau + \log \sigma(z_i)^{-1}))$$

bit operations. Thus, for the total cost, we obtain the bound

$$\tilde{O}(n^3\tau + n^2 \sum_{i=1}^n \log \sigma(z_i)^{-1}) = \tilde{O}(n^3\tau)$$

since $\sum_{i=1}^n \log \sigma(z_i)^{-1} = O(n\tau(\log n\tau)) = \tilde{O}(n\tau)$ according to Lemma 19 in [28].

3. Algorithm and Analysis

3.1. The Algorithm

We first present our new algorithm denoted DSC². For pseudo-code, we refer to the Appendix.

DSC² maintains a list $\mathcal{A}$ of active intervals I with corresponding integers $N_I = 2^{n_I}$, $n_I \in \mathbb{N}_{\geq 1}$, and a list $\mathcal{O}$ of isolating intervals, where we initially set $\mathcal{A} := \{(\mathcal{I}_0, 4)\} := \{((-2^{\tau+1}, 2^{\tau+1}), 4)\}$ and $\mathcal{O} := \emptyset$. For $(I, N_I) \in \mathcal{A}$, $I = (a, b)$, we proceed as follows: We remove I from $\mathcal{A}$ and compute the number $v := \text{var}(f, I)$ of sign variations for f on I .

- (1) If $v = 0$, we do nothing (i.e., I is discarded).
- (2) If $v = 1$, then I isolates a real root of f . Thus, we add I to the list $\mathcal{O}$ of isolating intervals.
- (3) For $v > 1$, we proceed as follows:

- (a) For $i = 1, 2$, let

$$B_1 := (a, a + \frac{w(I)}{N_I}) \quad \text{and} \quad B_2 := (b - \frac{w(I)}{N_I}, b) \quad (3.1)$$

be the left- and rightmost interval of size $w(I)/N_I$ contained in I , respectively. We compute $v_i := \text{var}(f, B_i)$: If one of the values v_1 or v_2 equals v , then the corresponding interval B_i (at most one of the two disjoint intervals B_i fulfills $\text{var}(f, B_i) = v$) contains all roots of f within I . Hence, we keep $I' := B_i$, discard $I \setminus I'$, and set $N_{I'} := N_I^2$. That is, $(I', N_{I'}) := (B_i, N_I^2)$ is added to $\mathcal{A}$.

- (b) If both values v_1 and v_2 differ from v , we compute

$$\lambda_1 := a - v \cdot \frac{f(a)}{f'(a)} \quad \text{and} \quad \lambda_2 := b - v \cdot \frac{f(b)}{f'(b)}. \quad (3.2)$$

If I contains a cluster $\mathcal{C}$ of v nearby roots and a (or b) has "reasonable" distance to $\mathcal{C}$, then λ_1 (or λ_2) constitutes a considerably better approximation of $\mathcal{C}$ than a (or b). We check whether this is actually the case: For $i = 1, 2$, we first compute the point $a + k_i \cdot \frac{w(I)}{4N_I}$, with $k_i \in \{2, \dots, 4N_I - 2\}$, which is closest to λ_i (if there exist two equally close points, we choose the one with smaller index). In more mathematical terms,

$$k_i := \min(\max(\lfloor 4N_I(\lambda_i - a) \rfloor, 2), 4N_I - 2). \quad (3.3)$$

Then, we define I'_i to be the interval of length $w(I)/N_I$ centered at the subdivision point $a + k_i \cdot w(I)/4N_I$, that is,

$$I'_i := (a + (k_i - 2) \cdot \frac{w(I)}{4N_I}, a + (k_i + 2) \cdot \frac{w(I)}{4N_I}) \subset I. \quad (3.4)$$

In particular, if $a + w(I)/4N_I \leq \lambda_i \leq b - w(I)/4N_I$, then I'_i contains λ_i , and λ_i has distance at least $w(I)/4N_I$ to both endpoints of I'_i . Now, we compute $v'_1 := \text{var}(f, I'_1)$ and $v'_2 := \text{var}(f, I'_2)$. If one of the two values v'_1 or v'_2 equals v , we keep the corresponding interval $I' := I'_i$ with $v'_i = v$ (if we count v sign variations for both intervals I'_1 and I'_2 , we just keep I'_1) and discard $I \setminus I'$. Finally, we add $(I', N_{I'}) := (I', N_I^2)$ to $\mathcal{A}$.

- (c) If all values v_1 , v_2 , v'_1 and v'_2 differ from v , then we consider this an indicator that there is either no cluster of v nearby roots or that such a cluster is not separated well enough from the remaining roots. Hence, in this situation, we fall back to bisection. That is, we split I into two equally sized intervals $I_1 = (a, m(I))$ and $I_2 = (m(I), b)$ and add $(I_1, \max(4, \sqrt{N}))$ and $(I_2, \max(4, \sqrt{N}))$ to $\mathcal{A}$. Finally, if $f(m(I)) = 0$, we also add $[m(I), m(I)]$ to $\mathcal{O}$.

Correctness and termination of DSC^2 are obvious because our starting interval $\mathcal{I}_0$ contains all real roots of f , we never discard intervals (or endpoints) that contain a root of f , and intervals are at least bisected in each iteration. In addition, we obtain the following bounds on the width of the intervals I and the corresponding numbers N_I produced by DSC^2 :

Lemma 1. For each interval I produced by DSC^2 , we have

$$2^{\tau+2} \geq w(I) \geq \sigma_f^3 \cdot 2^{-2(\tau+2)} = 2^{-\tilde{O}(n\tau)} \text{ and } 4 \leq N_I \leq 2^{2(\tau+2)} \cdot \sigma_f^{-2} = 2^{\tilde{O}(n\tau)}.$$

In particular, for $I \in T_i$ (see Section 2 for the definition of the subtree $T_i \subset T_{\text{DSC}^2}$), we have

$$2^{\tau+2} \geq w(I) \geq w(J_i) \text{ and } 4 \leq N_I \leq 2^{2(\tau+2)} \cdot w(J_i)^{-2},$$

with J_i the special node corresponding to T_i .

Proof. The inequalities $2^{\tau+2} \geq w(I)$ and $N_I \geq 4$ are trivial. For $N_I > 4$, there must exist an interval $J \supset I$ with $N_J = \sqrt{N_I}$, and J was replaced by an interval $J' \supseteq I$ of size $w(J)/N_J$. Since J is non-terminal, J' is also non-terminal because $\text{var}(f, J') = \text{var}(f, J) > 1$. Thus, $\sigma_f \leq w(J') = w(J)/N_J \leq 2^{\tau+2}/\sqrt{N_I}$. This shows the upper bound for N_I . For the lower bound for $w(I)$, we consider the parent interval J of I . Since J is non-terminal, we have $w(J) \geq \sigma_f$, and thus $w(I) \geq w(J)/N_J \geq \sigma_f \cdot \sigma_f^2 \cdot 2^{-2(\tau+2)}$. For $I \in T_i$, the bounds for $w(I)$ are trivial, and, in completely similar manner as above, we conclude that $2^{\tau+2}/\sqrt{N_I} \geq w(J_i)$. $\square$

Throughout the following considerations, we call a subdivision step from I to $I' \subset I$ *quadratic* if $w(I') = w(I)/N_I$, and we call a subdivision step *linear* if I is split into two equally sized intervals I_1 and I_2 . In a quadratic step, the integer N_I is squared whereas, in a linear step, $N_{I'} := \max(4, \sqrt{N_I})$ for each of the subintervals $I' = I_{1/2}$.

3.2. Analysis of the Recursion Tree

In this section, we prove that the size of each of the subtrees $T_i \subset T_{\text{DSC}^2}$ as defined in Section 2 is bounded by $O(\log n + \log \tau)$. We first have to investigate into the following technical lemmata:

Lemma 2. Let $w, w' \in \mathbb{R}^+$ be two positive reals with $w > w'$, and let $m \in \mathbb{N}_{\geq 1}$ be a positive integer. The sequence $(s_i)_{i \in \mathbb{N}_{\geq 1}} := ((x_i, n_i))_{i \in \mathbb{N}_{\geq 1}}$ is recursively defined as follows: $s_1 = (x_1, n_1) = (w, m)$, and

$$s_i = (x_i, n_i) := \begin{cases} \left(\frac{x_{i-1}}{N_{i-1}}, n_{i-1} + 1 \right), & \text{if } \frac{x_{i-1}}{N_{i-1}} \geq w' \\ \left(\frac{x_{i-1}}{2}, \max(1, n_{i-1} - 1) \right), & \text{if } \frac{x_{i-1}}{N_{i-1}} < w', \end{cases}$$

where $N_i := 2^{2^{n_i}}$ and $i \geq 2$. Then, the smallest index i_0 with $x_{i_0} \leq w'$ is upper bounded by $8(n_1 + \log \log \max(4, \frac{w}{w'}))$.

Proof. Throughout the following consideration, we call an index i *strong* (**S**) if $x_i/N_i \geq w'$ and *weak* (**W**), otherwise. If $w/4 < w'$, then each $i \geq 1$ is weak, and thus $i_0 \leq 3$. For $w/4 \geq w'$, let $k \in \mathbb{N}_{\geq 1}$ be the unique integer with

$$2^{-2^{k+1}} < w'/w \leq 2^{-2^k}.$$

Then, $k \leq \log \log \frac{w}{w'}$, and since $x_i \leq x_{i-1}/2$ for all i , there must exist an index i which is weak. Let k' denote the smallest weak index.

Claim 1: $k' \leq k + 1$

Assume otherwise, then the indices 1 to k are all strong. Hence,

$$x_{k+1} = w \cdot 2^{-(2^m + 2^{m+1} + \dots + 2^{m+k-1})} = w \cdot 2^{-2^m(2^0 + 2^1 + \dots + 2^{k-1})} = w \cdot 2^{-2^m(2^k - 1)} \leq 4w \cdot 2^{-2^{k+1}} < 4w',$$

and $n_{k+1} > 1$. It follows that $k + 1$ is weak, a contradiction.

Let us now consider the subsequence $\mathcal{S} = k', k' + 1, \dots, i_0 - 3$:

Claim 2: $\mathcal{S}$ contains no subsequence of type ...SS... or ...SWSWS...

If there exists a weak index i and two strong indices $i + 1$ and $i + 2$, then $x_i/N_i > x_{i+2}/N_i \geq x_{i+2}/N_{i+2} \geq w'$ contradicting the fact that $x_i/N_i < w'$. Since $\mathcal{S}$ starts with a weak index, the first part of our claim follows. For the second part, assume that $i, i + 2$ and $i + 4$ are strong. Then, $i + 1$ and $i + 3$ are weak, and thus

$$w' \leq x_{i+4}/N_{i+4} < x_{i+2}/(N_{i+2} \cdot N_{i+4}) < x_i/(N_i \cdot N_{i+2} \cdot N_{i+4}) = x_i/N_i^3 = x_{i+1}/N_{i+1}$$

contradicting the fact that $i + 1$ is weak.

Claim 3: If i is weak and $i < i_0$, then $n_i \geq 2$.

Namely, if i is weak and $n_i = 1$, then $x_i/4 = x_i/N_i < w'$, and thus $x_{i_0-1} < w'$ which contradicts the definition of i_0 .

We now partition the sequence $\mathcal{S}$ into maximal subsequences $\mathcal{S}_1, \mathcal{S}_2, \dots, \mathcal{S}_r$ such that each $\mathcal{S}_j$, $j = 1, \dots, r$, contains no two consecutive weak elements. Then, according to our above results, each $\mathcal{S}_j$, with $j < r$, is of type **W**, **WSW**, or **WSWSW**. The last subsequence $\mathcal{S}_r$ (with last index $i_0 - 3$) is of type **W**, **WS**, **WSW**, **WSWS**, or **WSWSW**. After each $\mathcal{S}_j$, with $j < r$, the number n_i decreases by one, and thus we must have $r \leq n_1 + k'$ since we start with $n_{k'} = n_1 + k' - 1$ and, in addition, $n_i \geq 2$ for all weak i . Since the length of each $\mathcal{S}_j$ is bounded by 5, it follows that

$$i_0 = i_0 - 3 + 3 \leq k' + 5r + 3 \leq 5(n_1 + k') + k' + 3 \leq 8(n_1 + k).$$

□

Lemma 3. Let $I = (a, b)$ an arbitrary interval, A_n the corresponding Obreshkoff area and L_n the Obreshkoff lens for I . Then:

(1) For $I' = (a', b') \subset I$ with $a \neq a'$ and $b \neq b'$, the Obreshkoff area A'_n for I' is completely contained within the lens L_n if

$$\min(|a - a'|, |b - b'|) > 8n^2 w(I').$$

In the latter situation, it holds that

$$|x - \xi| > \frac{1}{4n} \cdot (\min(|a - a'|, |b - b'|) - 8nw(I'))$$

for all $x \notin L_n$ and all $\xi \in A'_n$.

(2) For $I' = (a', b')$ with $I' \cap I = \emptyset$, the Obreshkoff area A'_n for I' does not intersect A_n if

$$\text{dist}(I, I') > 4n^2 \cdot \min(w(I), w(I')),$$

where $\text{dist}(I, I')$ denotes the distance between the two intervals I and I' .




Fig. 3.1. On the left figure, c denotes the topmost point of the Obreshkoff lens L_n for $I = (a, b)$. If $|a - a'| \leq w(I)/2$, then the distance from a' to the boundary of L_n is bounded by the distance δ from a' to $\overline{ac}$. The radius r' of the Obreshkoff discs $\underline{C}'_n$ and $\overline{C}'_n$ for $I' = (a', b')$ is bounded by $n \cdot w(I')$ due to the extended Sine Theorem. The right figure shows the Obreshkoff areas for the intervals I and I' , respectively.

Proof. (1) In a first step, we compute the radius r' of the Obreshkoff discs $\underline{C}'_n$ and $\overline{C}'_n$ for the interval $I' = (a', b')$: A point ξ on the boundary of the Obreshkoff area A'_n (except a' and b') sees I' under an angle $\gamma = \pi/(n+2)$; see Figure 2.1 and 3.1. Hence, from the extended Sine Theorem, it follows that

$$r' = \frac{w(I')}{2 \sin(\gamma)} = \frac{w(I')}{2 \sin(\pi/(n+2))} < \frac{(n+2)w(I')}{\pi} < n \cdot w(I')$$

since $\sin x > x/2$ for all $x \in (0, \pi/4]$. In particular, each point z within the Obreshkoff area A'_n has distance at most $2r' < 2n \cdot w(I')$ from any point within I . W.l.o.g., we assume that $|a - a'| \leq |b - b'|$. Then, the distance from a' to the boundary of the Obreshkoff lens L_n for I is bounded by the distance δ from a' to the line $\overline{ac}$, where c denotes the topmost point of L_n . Since $\overline{ac}$ intersects the x -axes in an angle of $\pi/(2(n+2))$, we have $\delta = |a - a'| \sin \pi/(2(n+2)) > |a - a'|/(4n)$. Thus, $A'_n \subset L_n$ if $|a - a'|/(4n) > 2n \cdot w(I')$ or $|a - a'| > 8n^2 w(I')$. For $|a - a'| > |b - b'|$, a similar argument shows that $A'_n \subset L_n$ if $|b - b'| > 8n^2 w(I')$. In addition, if $\min(|a - a'|, |b - b'|) > 8n^2 w(I')$, then each point $\xi \in A'_n$ has distance at least $\delta - 2nw(I') > \min(|a - a'|, |b - b'|)/(4n) - 2nw(I')$ from any point located outside of L_n .

(2) W.l.o.g., we can assume that $w(I') \leq w(I)$ and $a' \geq b$. Let L be the line passing through b which intersects the x -axes in an angle of $\pi/(n+2)$. Then, the upper part of the Obreshkoff area A_n lies completely on one side of this line. Now, if A'_n lies completely on the other side of L , then, by symmetry, A_n and A'_n do not share a common point. We have already argued that A'_n is contained within the disc of radius $2nw(I')$ centered at a' . Hence, if the distance $\delta' := \text{dist}(a', L)$ from a' to L is larger than $2nw(I')$, then $A_n \cap A'_n = \emptyset$. We have $\delta' = |a' - b| \sin(\pi/(n+2)) > |a' - b|/2n = \text{dist}(I, I')/2n$, and thus our claim follows. $\square$

We now turn to the analysis of the subtrees $T_i \subset T_{\text{DSC}^2}$ as defined in Section 2. There, we have already argued that each T_i constitutes a chain of intervals $I_1 = (a_1, b_1) \supset I_2 = (a_2, b_2) \supset \dots \supset I_s = (a_s, b_s)$ “connecting” the special node J_i with the special node J_k of minimal length that contains J_i . In the following Theorem, we will show that, for all but $O(\log n)$ many j , the sequence $(w(I_j), n_{I_j}) = (w(I_j), \log \log N_{I_j})$ behaves similar to the sequence (x_j, n_j) as defined in Lemma 2. As a result, we obtain the following bound on $|T_i|$:

Theorem 4. For each special node J_i , the corresponding subtree $T_i \subset T_{\text{DSC}^2}$ has size

$$|T_i| = O(\log n + \log \tau).$$

Proof. We first consider the special case where $a_1 = a_2 = \dots = a_s$, that is, in each subdivision step, the leftmost interval has been chosen. Since $v = \text{var}(I_1) = \dots = \text{var}(I_s)$, Theorem 2 implies that $\text{var}(f, I) = v$ for each interval I with $I_s \subset I \subset I_1$. In particular, if $w(I_j)/N_{I_j} \geq w(I_s)$, we count v sign variations for the interval $B_1 = (a_j, a_j + w(I_j)/N_{I_j}) = (a_s, a_s + w(I_j)/N_{I_j})$ as defined in (3.1).

Thus, the subdivision step from I_j to I_{j+1} is quadratic in this case. Then, for $j = 1, \dots, s-1$, the sequence $(w(I_j), n_{I_j})$ coincides with the sequence (x_j, n_j) as defined in Lemma 2, where $w := w(I_1)$, $w' := w(I_s)$ and $n_1 = m := n_{I_1}$. Namely, if $w(I_j)/N_{I_j} \geq w'$, we have $w(I_{j+1}) = w(I_j)/N_{I_j}$ and $n_{I_{j+1}} = 1 + n_{I_j}$, and, otherwise, we have $w(I_{j+1}) = w(I_j)/2$ and $n_{I_{j+1}} = \max(1, n_{I_j} - 1)$. Hence, according to Lemma 2, it follows that s is bounded by

$$8(n_{I_1} + \log \log \max(4, w(I_1)/w(I_s))) = O(\log n + \log \tau),$$

where we used the bounds on n_{I_1} and $w(I_s)$ from Lemma 1. An analogous argument shows the same bound for s in the case where $b_1 = b_2 = \dots = b_s$.

We now turn to the more general case, where $a_1 \neq a_s$ and $b_1 \neq b_s$: Let $s_1 \in \{1, \dots, s\}$ be the smallest index with $a_{s_1} \neq a_1$ and $b_{s_1} \neq b_1$. Then, due to the above argument, s_1 is bounded by $O(\log n + \log \tau)$. Furthermore, $\min(|a_1 - a_{s_1}|, |b_1 - b_{s_1}|) \geq w(I_{s_1})/4$, and thus

$$\begin{aligned} \min(|a_1 - a_j|, |b_1 - b_j|) &\geq 2^{j-s_1-2} w(I_j) \\ \Rightarrow \frac{1}{4n} \cdot (\min(|a_1 - a_j|, |b_1 - b_j|) - 8nw(I_j)) &\geq w(I_j) \left(\frac{2^{j-s_1-4}}{n} - 8n \right) \text{ for all } j \geq s_1. \end{aligned}$$

Hence, with $s_2 := s_1 + \lceil \log(16n^3) \rceil + 4 = O(s_1) + O(\log n)$, this yields

$$\frac{1}{4n} \cdot (\min(|a_1 - a_j|, |b_1 - b_j|) - 8nw(I_j)) \geq 8n^2 w(I_j) \text{ for all } j \geq s_2.$$

Then, from Theorem 1 and Lemma 3, we conclude that, for $j \geq s_2$, the Obreshkoff area $A_n^{(j)}$ for I_j contains exactly v roots $z_1, \dots, z_v$ of f because the Obreshkoff lens $L_n^{(1)}$ for I_1 contains at most v roots and $A_n^{(j)} \subset L_n^{(1)}$ contains at least v roots. In particular, the Obreshkoff area $A_n^{(s)}$ for I_s must contain $z_1, \dots, z_v$. In the proof of Lemma 3, we have already argued that each point within $A_n^{(s)}$ has distance less than $2nw(I_s)$ from any point within I_s , and thus

$$|x - z_i| < 2nw(I_s), \text{ for all } i = 1, \dots, v \text{ and all } x \in I_s. \quad (3.5)$$

The remaining roots $z_{v+1}, \dots, z_n$ of f are located outside the Obreshkoff lens $L_n^{(1)}$ for I_1 , and thus their distance to an arbitrary point within I_j is larger than $8n^2 w(I_{s_2})$. Namely, according to Lemma 3, the distance from any of the roots $z_{v+1}, \dots, z_n$ to an arbitrary point within I_{s_2} is lower bounded by $8n^2 w(I_{s_2})$ and I_{s_2} contains I_j . The following consideration further shows the existence of an $s_3 = s_2 + O(\log n + \log \tau)$ such that $w(I_j) \leq w(I_{s_2})/N_{I_j}$ for all $j \geq s_3$, and thus

$$|x - z_i| > 8n^2 N_{I_j} w(I_j) \text{ for all } i = v+1, \dots, n, j \geq s_3, \text{ and all } x \in I_j: \quad (3.6)$$

Due to Lemma 1, we have $N_{I_j} \leq N_{\max} := \lceil 2^{2(\tau+2)} / \sigma_f \rceil = 2^{\tilde{O}(n\tau)}$ for all j . Thus, if the sequence $I_{s_2}, I_{s_2+1}, \dots$ starts with more than $m_{\max} := \log \log N_{\max} + 1 = O(\log n + \log \tau)$ consecutive linear subdivision steps, then $N_{I_{j'}} = 4$ and $w(I_{j'}) \leq w(I_{s_2})/4 = w(I_{s_1})/N_{I_{j'}}$ for some $j' \leq s_2 + \log \log m_{\max}$. Otherwise, there exists a j' with $s_2 \leq j' \leq s_2 + m_{\max}$ such that the step from $I_{j'}$ to $I_{j'+1}$ is quadratic. Since the length of a sequence of consecutive quadratic subdivision steps is also bounded by $m_{\max}$, there must exist a j'' with $j' + 1 \leq j'' \leq j' + m_{\max} + 1$ such that the step from $I_{j''-1}$ to $I_{j''}$ is quadratic, whereas the step from $I_{j''}$ to $I_{j''+1}$ is linear. Then, $N_{I_{j''+1}} = \sqrt{N_{I_{j''}}} = N_{I_{j''-1}}$ and

$$w(I_{j''+1}) = w(I_{j''})/2 = w(I_{j''-1})/(2N_{I_{j''-1}}) < w(I_{s_2})/N_{I_{j''+1}}.$$

Hence, in both cases, we have shown that there exists an $s_3 \leq s_2 + 2m_{\max} + 1 = O(\log n + \log \tau)$ with $w(I_{s_3}) \leq w(I_{s_2})/N_{I_{s_3}}$. Then, by induction, it follows that $w(I_j) \leq w(I_{s_2})/N_{I_j}$ for all $j \geq s_3$ which shows (3.6).

We are now ready to show that the subdivision step from I_j to I_{j+1} is quadratic if $j \geq s_3$ and $w(I_j) \geq 68nN_{I_j}w(I_s)$: Namely, if the latter two inequalities hold, then one of the endpoints (w.l.o.g., we assume this point to be a_j) of I_j has distance at least $w(I_j)/2 \geq 34nN_{I_j}w(I_s)$ from a_s . Thus, the distance from a_j to any of the roots $z_1, \dots, z_v$ is larger than $34nN_{I_j}w(I_s) - 2nw(I_s) \geq 32nN_{I_j}w(I_s)$. In addition, we have $|a_j - z_i| > 8n^2N_{I_j}w(I_j)$ for all $i > v + 1$ due to (3.6). Thus,

$$\begin{aligned} \left| \frac{1}{v} \cdot \frac{(a_j - a_s)f'(a_j)}{f(a_j)} \right| &= \left| \frac{1}{v} \sum_{i=1}^v \frac{a_j - a_s}{a_j - z_i} + \frac{1}{v} \sum_{i=v+1}^n \frac{a_j - a_s}{a_j - z_i} \right| = \left| 1 + \frac{1}{v} \sum_{i=1}^v \frac{z_i - a_s}{a_j - z_i} + \frac{1}{v} \sum_{i=v+1}^n \frac{a_j - a_s}{a_j - z_i} \right| \\ &\leq 1 + \frac{1}{v} \sum_{i=1}^v \frac{|z_i - a_s|}{|a_j - z_i|} + \frac{1}{v} \sum_{i=v+1}^n \frac{|a_j - a_s|}{|a_j - z_i|} < 1 + \frac{2nw(I_s)}{32nN_{I_j}w(I_s)} + n \cdot \frac{w(I_j)}{8n^2N_{I_j}w(I_j)} \\ &\leq 1 + \frac{1}{16N_{I_j}} + \frac{1}{8nN_{I_j}} \leq 1 + \frac{1}{8N_{I_j}}, \end{aligned}$$

where we used that $f'(a)/f(a) = \sum_{i=1}^n (a - z_i)^{-1}$ for all $a \in \mathbb{C}$ with $f(a) \neq 0$. In completely analogous manner, we show that

$$\left| \frac{1}{v} \cdot \frac{(a_j - a_s)f'(a_j)}{f(a_j)} \right| > 1 - \frac{1}{8N_{I_j}}.$$

This yields the existence of an $\varepsilon \in \mathbb{R}$ with $|\varepsilon| < 1/(8N_j) \leq 1/32$ and $\frac{1}{v} \cdot \frac{(a_j - a_s)f'(a_j)}{f(a_j)} = 1 + \varepsilon$. We can now derive the following bound on the distance between the approximation obtained by the Newton iteration and a_s :

$$\begin{aligned} \left| a_s - (a_j - v \cdot \frac{f(a_j)}{f'(a_j)}) \right| &= |a_s - a_j| \cdot \left| 1 - \frac{1}{\frac{1}{v} \cdot \frac{(a_j - a_s)f'(a_j)}{f(a_j)}} \right| = |a_s - a_j| \cdot \left| 1 - \frac{1}{1 + \varepsilon} \right| \\ &= \varepsilon \cdot |a_s - a_j| \cdot \left| \frac{1}{1 + \varepsilon} \right| < \frac{4|a_s - a_j|}{33N_{I_j}} \leq \frac{w(I_j)}{7N_{I_j}} \end{aligned} \quad (3.7)$$

If $a_s \geq b_j - w(I_j)/N_{I_j}$, then $(a_s, b_s) \subset B_2 = (b_j - w(I_j)/N_{I_j}, b_j)$ (cf. Step 3 (a), (3.1) in our algorithm for the definition of B_2), and thus $\text{var}(f, B_2) = v$. Hence, in this case, we keep $I_{j+1} = B_2$ which has size $w(I_j)/N_{I_j}$. If $a_s < b_j - w(I_j)/N_{I_j}$, then according to (3.7) we have

$$a_j - v \cdot \frac{f(a_j)}{f'(a_j)} \in (a_s - \frac{w(I_j)}{7N_{I_j}}, a_s + \frac{w(I_j)}{7N_{I_j}}) \subset (a_j + \frac{w(I_j)}{4N_{I_j}}, b_j - \frac{w(I_j)}{4N_{I_j}}).$$

It follows that the interval I'_1 as defined in Step 3 (b), (3.4) of our algorithm contains (a_s, b_s) , and thus $\text{var}(f, I'_1) = v$. This shows that the subdivision step from I_j to I_{j+1} is quadratic.

We now consider the sequence $(w(I_{s_3+i}), n_{I_{s_3+i}})$, for $i = 1, \dots, i'$, where i' is defined as the largest index with $w(I_{s_3+i'}) \geq 68nw(I_s)$. Then, our above argument implies that the sequence $(w(I_{s_3+i}), n_{I_{s_3+i}})_{1 \leq i \leq i'-1}$ coincides with the sequence $(x_i, n_i)_{1 \leq i \leq i'-1}$ as defined in Lemma 2, where $n_1 = m = n_{I_{s_3+1}}$ and $w' := 68nw(I_s)$. Namely, if $w(I_{s_3+i})/N_{I_{s_3+i}} \geq w'$, then $68nw(I_s) \leq w(I_{s_3+i+1}) = w(I_{s_3+i})/N_{I_{s_3+i}}$ and $n_{I_{s_3+i+1}} = 1 + n_{I_{s_3+i}}$, whereas, for $w(I_{s_3+i})/N_{I_{s_3+i}} < w'$, we have $w(I_{s_3+i+1}) = w(I_{s_3+i})/2$ and $n_{I_{s_3+i+1}} = \max(n_{I_{s_3+i}} - 1, 1)$. It follows that i' is bounded by $8(n_1 + \log \log \max(4, w(I_{s_3+1})/w')) = O(\log n + \log \tau)$. Hence, there exists an $s_4 = s_3 + i' + 1 = O(\log n +$

$\log \tau$) with $w(I_j) < 68nw(I_s)$ for all $j \geq s_4$. Finally, this shows that s is upper bounded by $s_4 + \log(68n) = O(\log n + \log \tau)$. $\square$

Combining the latter theorem and (2.3) immediately yields the following result on the size of the induced recursion tree:

Theorem 5. For a polynomial f of degree n with integer coefficients of bitsize τ , the algorithm DSC^2 induces a recursion tree T_{DSC^2} of size

$$|T_{\text{DSC}^2}| = \text{var}(f, I_0) \cdot O(\log n + \log \tau) = O(n \cdot (\log n + \log \tau)),$$

where $I_0 := (-2^{\tau+1}, 2^{\tau+1})$ denotes the initial interval known to contain all real roots of f .

3.3. Bit Complexity Analysis

We now derive an upper bound for the number of bit operations that are needed to determine isolating intervals for the real roots of f . We will show that, in each iteration, the costs are dominated (up to a constant factor) by the computation of the polynomial $f_I = (x+1)^n \cdot f((ax+b)/(x+1))$ as defined in (2.2). The costs for this step mainly depend on the bitsize of the endpoints, and thus on the length of the interval $I = (a, b)$. The following Lemma provides a lower bound on the width of the special nodes J_i , and thus also for the nodes $I \in T_i$, in terms of the separations of the roots $z_1, \dots, z_n$:

Lemma 6. For a polynomial f as defined in (2.1), we can order the roots of $z_1, \dots, z_n$ of f in way such that

$$w(J_i) > \frac{\sigma(z_i)}{4} \cdot n^{-5-2\log n},$$

where $i = 1, \dots, n'$ and $J_1, \dots, J_{n'}$ denote the special nodes in the recursion tree T_{DSC^2} . In addition, the endpoints of an arbitrary interval $I \in T(i) \cup \{J_i\}$ are dyadic numbers which can be represented by $O(\tau + \log \sigma_i^{-1} + \log^2 n)$ bits.

Proof. We first order the roots of f with respect to their separations, that is, $\sigma(z_1) \geq \dots \geq \sigma(z_n)$. For a fixed $\sigma \in \mathbb{R}^+$, let k be defined such that $\sigma(z_{n-k+1}) < \sigma \leq \sigma(z_{n-k})$, that is, exactly the k roots $z_{n-k+1}, \dots, z_n$ have separation less than σ . We further denote $I_1 := J_{i_1}, \dots, I_m := J_{i_m}$ the special nodes such that

$$w(I_l) < w_{\min} := \frac{\sigma}{4} \cdot n^{-5-2\log n} \text{ for all } l = 1, \dots, m,$$

and each special node $J_i \in T_{\text{DSC}^2}$ which contains I_l has width $w(J_i) \geq w_{\min}$. In addition, we denote $v_l := \text{var}(f, I_l) \geq 2$ the number of sign variations that we count for I_l . Since the intervals I_l are disjoint, we have $v_1 + \dots + v_m \leq n$, and thus $m \leq n/2$. According to Theorem 1, v_l is a lower bound for the number of roots within the Obreshkoff area $A_n^{(l)}$ for I_l . Furthermore, in Lemma 3, we have proven that any two points within $A_n^{(l)}$ have distance less than $4nw(I_l) < \sigma$, and thus each root contained in $A_n^{(l)}$ must be one of the k roots $z_{n-k+1}, \dots, z_n$. Let S_l be the set of all roots which are contained in $A_n^{(l)}$.

Let us first consider the case, where the Obreshkoff areas $A_n^{(l)}$ are pairwise disjoint. Then, the subsets $S_l \subset A_n^{(l)}$ are also pairwise disjoint, and thus

$$v_1 + \dots + v_s \leq |S_1| + \dots + |S_m| \leq k.$$

We now turn to the more general situation, where some of the $A_n^{(l)}$ may overlap. If this happens, then it is possible that, in total, some of the roots contained in these areas are counted more than once. Hence, we propose the following approach: In a first step (see the subsequent construction), starting with the intervals $I_1, \dots, I_m$, we iteratively merge intervals whose corresponding Obreshkoff areas overlap until we finally obtain intervals $I'_1, \dots, I'_{m'}$ such that

- the intervals $I_1, \dots, I_m$ are covered by $I'_1, \dots, I'_{m'}$,
- each I'_l has width $w(I'_l) < w_{\min} \cdot n^{4+2\log n}$, and
- the Obreshkoff areas A_n for I'_l are pairwise disjoint.

Then, using the same argument as above yields $v_1 + \dots + v_m \leq v' := \sum_{l=1}^{m'} \text{var}(f, I'_l) \leq k$, where the latter inequality follows from the fact that the Obreshkoff area A_n for each I'_l contains only roots with separation less than $4nw(I'_l) < 4nw_{\min} \cdot n^{4+2\log n} < \sigma(z_{n-k})$. It remains to show how to construct the intervals I'_l : We start with a list $\mathcal{A}$ of active intervals, where we initially set $\mathcal{A} := \{I_1, \dots, I_m\}$. In each iteration, we pick two intervals $I = (a, b)$ and $I' = (a', b')$ from $\mathcal{A}$ whose corresponding Obreshkoff areas overlap. Then, we remove I, I' , and all intervals $J \in \mathcal{A}$ in between I and I' . Finally, we add the smallest interval K into $\mathcal{A}$ which contains I and I' (i.e., $K = (\min(a, a'), \max(b, b'))$). We proceed in this way until we obtain intervals $I'_1, \dots, I'_{m'}$ such that the corresponding Obreshkoff areas do not overlap. It remains to show that the length of each of the so-obtained intervals is bounded by $w_{\min} \cdot n^{4+2\log n}$: At any stage, an interval J obtained in the above iteration, covers a certain number s of intervals from $I_1, \dots, I_m$. By induction on s , we prove that

$$w(J) < w_{\min} \cdot s^4 n^{2\log s}, \text{ and thus } w(J) \leq w_{\min} \cdot n^{2\log n + 4}. \quad (3.8)$$

If $s = 1$, then trivially $J = I_l$ for some l which proves the claim for $s = 1$. An interval J covering $s + 1$ intervals from $I_1, \dots, I_m$ is obtained by merging two intervals I and I' which cover s_1 and s_2 intervals, respectively, where $s_1 + s_2 \leq s + 1$ and, w.l.o.g., $1 \leq s_1 \leq s_2$. From Lemma 3, it follows that the distance between I and I' is bounded by $4n^2 \cdot \min(w(I'), w(I))$ because the corresponding Obreshkoff areas overlap. Hence, J has width

$$\begin{aligned} w(J) &\leq w(I) + w(I') + 4n^2 \cdot \min(w(I'), w(I)) < w_{\min} \cdot (s_2^4 n^{2\log s_2} + (4n^2 + 1)s_1^4 n^{2\log s_1}) \\ &< w_{\min} \cdot (s_2^4 n^{2\log s_2} + 8s_1^4 n^{2\log s_1 + 2}) = w_{\min} \cdot (s_2^4 n^{2\log s_2} + 8s_1^4 n^{2\log 2s_1}) \end{aligned}$$

If $2s_1 \geq s_2$, then

$$s_2^4 n^{2\log s_2} + 8s_1^4 n^{2\log 2s_1} \leq (8s_1^4 + s_2^4) n^{2\log 2s_1} \leq (s_1 + s_2)^4 n^{2\log(s_1 + s_2)}.$$

Otherwise, we have

$$s_2^4 n^{2\log s_2} + 8s_1^4 n^{2\log 2s_1} \leq (8s_1^4 + s_2^4) n^{2\log s_2} < (s_1 + s_2)^4 n^{2\log(s_1 + s_2)},$$

and thus (3.8) follows.

Hence, we count at most k sign variations for the intervals I_l , in total. Now, the same argument as used in Section 2 to show that there exists at most $n' = \text{var}(f, [-2^{\tau+1}, 2^{\tau+1}])$ special nodes also shows that the number of all special nodes J with $w(J) \leq \sigma$ is bounded by k . Namely, we start with special nodes $I_1, \dots, I_m$ with $\sum_{l=1}^m \text{var}(f, I_l) \leq k$, and whenever a special node is subdivided, the value $\mu := \sum_l \text{var}(f, I) - \#\{I : \text{var}(f, I) \geq 2\}$ decreases by at least one. For the result on the width of the intervals I_j , we consider integers $k_1, \dots, k_s$, with $k_1 < k_2 < \dots < k_s$, such that

$$\sigma(z_1) = \dots = \sigma(z_{n-k_s}) > \sigma(z_{n-k_s+1}) = \dots = \sigma(z_{n-k_{s-1}}) > \sigma(z_{n-k_{s-1}+1}) = \dots = \sigma(z_n).$$

There exist at most k_j special nodes J_i with $w(J_i) < \frac{\sigma(z_{n-k_j})}{4} n^{-4-2\log n}$, and there cannot exist a special node J_i with $w(J_i) < \sigma_f = \sigma(z_n)$. In addition, the number of special nodes with

$w(J_i) < \sigma(z_{n-k_1})n^{-4-2\log n}/4$ is bounded by k_1 . Then, by induction, it follows that $w(J_i) < \sigma(z_i)n^{-4-2\log n}/4$ because the J_i are ordered with respect to their length.

For the bit complexity of the endpoints of an interval $I = (a, b) \in T(i) \cap \{J_i\}$, we remark that, due to our construction, a and b are both dyadic numbers with modulus bounded by $2^{\tau+1}$. In addition, the denominator of a and b in its dyadic representation is bounded by $\max(1, \log w(I)^{-1})$. Hence, we can represent both endpoints of I with $O(\tau + \log \sigma_i^{-1} + \log^2 n)$ many bits. $\square$

We will now derive our final result on the bit complexity of DSC²: In each step of the algorithm, we have to compute the polynomial $f_I(x) = (x+1)^n \cdot f((ax+b)/(x+1))$, where $I = (a, b)$ is the interval that is actually processed. The latter computation decomposes into computing $f_I^* := f(a + (b-a)x)$, reversing the coefficients, and then applying a Taylor shift by 1 (i.e., $x \mapsto x+1$). For the computation of f_I^* , we first shift f by a , and then scale by a factor $b-a = w(I)$ which is a power of two. Using asymptotically fast Taylor shift [11,32], the shift $x \mapsto x+a$ (i.e., the computation of $f(x+a)$) demands for $\tilde{O}(n^2(\tau + \log w(I)^{-1}))$ bit operations. The scaling $x \mapsto (b-a) \cdot x$ is achieved by just shifting the i -th coefficient of $f(x+a)$ by $i \cdot \log(b-a)^{-1} = i \cdot \log w(I)^{-1}$ many bits. Then, the resulting polynomial f_I^* has coefficients of modulus $2^{O(n\tau)}$, and the denominators of their dyadic representations are bounded by $2^{O(n(\tau + \log w(I)^{-1}))}$. Hence, reversing the coefficients of f_I^* , and then applying a Taylor shift by 1, demands for $\tilde{O}(n^2(\tau + \log w(I)^{-1}))$ bit operations. In summary, the costs for computing f_I are bounded by $\tilde{O}(n^2(\tau + \log w(I)^{-1}))$. The same bound further applies to the computation of λ_1 and λ_2 in Step 3 (b), (3.2) of our algorithm because, in this step, we have to evaluate a polynomial of degree n and bitsize τ at a $(\tau + \log w(I)^{-1})$ -bit number. If I is non-terminal, then we also have to compute the number of sign variations for the intervals B_1, B_2, I'_1 and I'_2 . The same argument as above also shows that we can do so using $\tilde{O}(n^2(\tau + \log N_I + \log w(I)^{-1}))$ bit operations.

For an interval $I \in T(i) \cup J_i$, we have

$$\log N_I + \log w(I)^{-1} = O(\log N_I + \log w(I_j)^{-1}) = O(\tau + \log \sigma(z_i)^{-1} + \log^2 n);$$

see Lemma 1 and 6. Thus, the total costs for all computations at I are bounded by $\tilde{O}(n^2(\tau + \sigma(z_i)^{-1}))$ bit operations. It remains to consider a terminal interval I which is one of the two children of a special node J_i . In this case, we only have to bound the cost for the computation of f_I because $\text{var}(f, I) \leq 1$. Since $w(I) = w(J_i)/2$, the latter computation needs $\tilde{O}(n^2(\tau + \log w(I_j)^{-1})) = \tilde{O}(n^2(\tau + \log \sigma(z_i)^{-1}))$ bit operations as well.

In Theorem 4, we have shown that $|T_i| = O(\log n + \log \tau)$ for all i , and thus the total costs for isolating the real roots of f are bounded by

$$O(\log n + \log \tau) \cdot \sum_{i=1}^{n'} \tilde{O}(n^2(\tau + \log \sigma(z_i)^{-1})) = \tilde{O}(n^3 \tau)$$

since $\sum_{i=1}^{n'} \log \sigma(z_i)^{-1} = O(n\tau + \sum_{i=1}^n \log \sigma(z_i)^{-1}) = \tilde{O}(n\tau)$ according to Lemma 19 in [28]. We fix this result:

Theorem 7. For a polynomial f of degree n with integer coefficients of modulus less than 2^τ , DSC² isolates the real roots of f using no more than $\tilde{O}(n^3 \tau)$ bit operations.

4. Conclusion

We introduced the first subdivision method to isolate the real roots of a polynomial which achieves the record bound $\tilde{O}(n^3\tau)$ for the bit complexity of this fundamental problem. In comparison to the asymptotically fast algorithms by Pan and Schönhage from the 80tes which compute all complex roots, the new approach is much simpler and can be considered practical to an extremely high degree. The algorithm is based on a novel subdivision technique which combines Descartes' Rule of Signs and Newton iteration. As a consequence, our algorithm shows quadratic convergence towards the roots in most steps.

So far, the approach only applies to polynomials with integer (or rational) coefficients. In [27], we showed how to modify a subdivision algorithm which uses exact computation in each step such that it also applies to polynomials f with arbitrary real coefficients that can be approximated to any specified error bound (bitstream coefficients). Following this approach, it seems reasonable to express the bit complexity in terms of the geometry of the roots $z_1, \dots, z_n$ of f . For instance, the modified version of the Descartes method from [28] isolates all real roots using $\tilde{O}(n(n\Gamma + \Sigma)^2)$ bit operations, where Γ constitutes a bound on the logarithm of the modulus of the roots and $\Sigma := \sum_{i=1}^n \log \sigma(z_i)^{-1}$. Due to the quadratic convergence achieved by the new algorithm, we expect a corresponding bitstream version to perform the same task by a considerably lower number of bit operations.

References

- [1] J. Abbott. Quadratic interval refinement for real roots. Poster presented at the 2006 Int. Symp. on Symb. and Alg. Comp. (ISSAC 2006).
- [2] A. G. Akritas and A. Strzeboński. A comparative study of two real root isolation methods. *Nonlinear Analysis:Modelling and Control*, 10(4):297–304, 2005.
- [3] A. Alesina and M. Galuzzi. A new proof of Vicent’s theorem. *L’Enseignement Mathématique*, 44:219–256, 1998.
- [4] M. Burr and F. Krahmer. Sqfreeeval: An (almost) optimal real-root isolation algorithm. *CoRR*, abs/1102.5266, 2011.
- [5] M. Burr, F. Krahmer, and C. Yap. Continuous amortization: A non-probabilistic adaptive analysis technique. *Electronic Colloquium on Computational Complexity (ECCC)*, 16:136, 2009.
- [6] M. Burr, V. Sharma, and C. Yap. Evaluation-based root isolation, Feb. 2009. In preparation.
- [7] G. E. Collins and A. G. Akritas. Polynomial real root isolation using Descartes’ rule of signs. In *ISSAC*, pages 272–275, 1976.
- [8] Z. Du, V. Sharma, and C. Yap. Amortized bounds for root isolation via Sturm sequences. In *SNC*, pages 113–130, 2007.
- [9] A. Eigenwillig. *Real Root Isolation for Exact and Approximate Polynomials using Descartes’ Rule of Signs*. PhD thesis, Universität des Saarlandes, May 2008.
- [10] A. Eigenwillig, V. Sharma, and C. Yap. Almost tight complexity bounds for the Descartes method. In *ISSAC*, pages 71–78, 2006.
- [11] J. Gerhard. Modular algorithms in symbolic summation and symbolic integration. *LNCS, Springer*, 3218, 2004.
- [12] X. Gourdon. *Combinatoire, Algorithmique et Géométrie des Polynômes*. Thèse, École polytechnique, 1996.
- [13] M. Hemmer, E. P. Tsigaridas, Z. Zafeirakopoulos, I. Z. Emiris, M. I. Karavelas, and B. Mourrain. Experimental evaluation and cross benchmarking of univariate real solvers. In *SNC*, pages 45–54, 2009.
- [14] J. Johnson. Algorithms for polynomial real root isolation. In B. Caviness and J. Johnson, editors, *Quantifier Elimination and Cylindrical Algebraic Decomposition*, Texts and monographs in Symbolic Computation, pages 269–299. Springer, 1998.
- [15] W. Krandick and K. Mehlhorn. New bounds for the Descartes method. *J. Symbolic Computation*, 41(1):49–66, 2006.
- [16] T. Lickteig and M.-F. Roy. Sylvester-Habicht sequences and fast Cauchy index computation. *J. of Symbolic Computation*, 31:315–341, 2001.
- [17] D. P. Mitchell. Robust ray intersection with interval arithmetic. In *Graphics Interface’90*, pages 68–74, 1990.
- [18] B. Mourrain, F. Rouillier, and M.-F. Roy. The Bernstein basis and real root isolation. In *Combinatorial and Computational Geometry*, pages 459–478, 2005.
- [19] N. Obrechhoff. *Zeros of Polynomials*. Marina Drinov, Sofia, 2003. Translation of the Bulgarian original.
- [20] N. Obrechhoff. Über die Wurzeln von algebraischen Gleichungen. *Jahresbericht der Deutschen Mathematiker-Vereinigung*, 33:52–64, 1925.
- [21] N. Obrechhoff. *Verteilung und Berechnung der Nullstellen reeller Polynome*. VEB Deutscher Verlag der Wissenschaften, 1963.
- [22] A. M. Ostrowski. Note on Vincent’s theorem. *Annals of Mathematics, Second Series*, 52(3):702–707, 1950. Reprinted in: Alexander Ostrowski, *Collected Mathematical Papers*, vol. 1, Birkhäuser Verlag, 1983, pp. 728–733.
- [23] V. Y. Pan. Sequential and parallel complexity of approximate evaluation of polynomial zeros. *Comput. Math. Applic.*, 14(8):591–622, 1987.
- [24] V. Y. Pan. Solving a polynomial equation: some history and recent progress. *SIAM Review*, 39(2):187–220, 1997.
- [25] D. Reischert. Asymptotically fast computation of subresultants. In *ISSAC 97*, pages 233–240, 1997. Maui, Hawaii.

- [26] F. Rouillier and P. Zimmermann. Efficient isolation of [a] polynomial's real roots. *J. Computational and Applied Mathematics*, 162:33–50, 2004.
- [27] M. Sagraloff. A general approach to isolating roots of a bitstream polynomial. *Mathematics in Computer Science, Special Issue on Algorithms and Complexity at the Interface of Mathematics and Computer Science*, 2011. to appear, see <http://www.mpi-inf.mpg.de/msagrало/bitstream10.pdf> for an online version.
- [28] M. Sagraloff. On the complexity of real root isolation. arXiv:1011.0344v2, 2011.
- [29] A. Schönhage. The fundamental theorem of algebra in terms of computational complexity, 1982. Manuscript, Department of Mathematics, University of Tübingen. Updated 2004.
- [30] V. Sharma. Complexity of real root isolation using continued fractions. *Theor. Computer Science*, 409(2), 2008. Also: proceedings ISSAC'07.
- [31] E. P. Tsigaridas and I. Z. Emiris. On the complexity of real root isolation using continued fractions. *Theor. Comput. Sci.*, 392(1-3):158–173, 2008.
- [32] J. von zur Gathen and J. Gerhard. Fast algorithms for Taylor shifts and certain difference equations. In *ISSAC '97: Proceedings of the 1997 international symposium on Symbolic and algebraic computation*, pages 40–47, New York, NY, USA, 1997. ACM.
- [33] C.-K. Yap and M. Sagraloff. A simple but exact and efficient algorithm for complex root isolation. In *ISSAC*, pages 353–360, 2011.

5. Appendix

Algorithm 1 DSC²

Require: polynomial $f = \sum_{0 \leq i \leq n} a_i x^i \in \mathbb{Z}[x]$ with integer coefficients a_i , $|a_i| < 2^\tau$ for all i .

Ensure: returns a list $\mathcal{O}$ of disjoint isolating intervals for all real roots of f

```

 $I_0 := (-2^{\tau+1}, 2^{\tau+1}); N_{I_0} := 4$ 
 $\mathcal{A} := \{ (I_0, N_{I_0}) \}; \mathcal{O} := \emptyset$  {list of active and isolating intervals}
repeat
   $(I, N_I)$  some element in  $\mathcal{A}$  with  $I = (a, b)$ ; delete  $(I, N_I)$  from  $\mathcal{A}$ 
   $v := \text{var}(f, I)$ 
  if  $v = 0$  then
    do nothing {I contains no root}
  else if  $v = 1$  then
    add  $I$  to  $\mathcal{O}$  {I isolates a real root}
  else if  $v > 1$  then
     $B_1 := (a, a + \frac{w(I)}{N_I}); B_2 := (b - \frac{w(I)}{N_I}, b)$ 
    if  $\text{var}(f, B_1) = v$  or  $\text{var}(f, B_2) = v$  then
      for the unique  $i \in \{1, 2\}$  with  $\text{var}(f, B_i) = v$ , add  $(B_i, N_{B_i}) := (B_i, N_I^2)$  to  $\mathcal{A}$ 
      {B_i contains all real roots within I }
    else
       $\lambda_1 := a - v \cdot \frac{f(a)}{f'(a)}; \lambda_2 := b - v \cdot \frac{f(b)}{f'(b)}$ 
      for  $i = 1, 2$ :
         $k_i := \min(\max(\lfloor 4N_I \cdot \frac{\lambda_i - a}{b - a} \rfloor, 2), 4N_I - 2); I'_i := (a + (k_i - 2) \cdot \frac{w(I)}{4N_I}, a + (k_i + 2) \cdot \frac{w(I)}{4N_I});$ 
         $\{m_k := a + k \cdot \frac{w(I)}{4N_I} \text{ is the } k\text{-th subdivision point when decomposing } I \text{ into } 4N_I \text{ equally}$ 
         $\text{sized intervals; } m_{k_i} \text{ is closest to } \lambda_i; I'_i \text{ has width } \frac{w(I)}{N_I} \text{ and is centered at } m_{k_i}\}$ 
      if  $\text{var}(f, I'_1) = v$  or  $\text{var}(f, I'_2) = v$  then
        choose the smallest  $i \in \{1, 2\}$  with  $\text{var}(f, I'_i) = v$  and add  $(I'_i, N_{I'_i}) := (I'_i, N_I^2)$  to  $\mathcal{A}$ 
        {If } \text{var}(f, I'_i) = v, \text{ then } I'_i \text{ contains all roots within } I \}
      else
        add  $(I_1, N_{I_1}) := ((a, m(I)), \max(4, \sqrt{N_I})), (I_2, N_{I_2}) := ((m(I), b), \max(4, \sqrt{N_I}))$  to  $\mathcal{A}$ 
        if  $f(m(I)) = 0$  then
          add  $[m(I), m(I)]$  to  $\mathcal{O}$ 
        end if
      end if
    end if
  end if
until  $\mathcal{A}$  is empty
return  $\mathcal{O}$ 

```
